

## VIII Covering Spaces

### A. Covering Spaces

recall, when we computed  $\pi_1(S^1)$  we used the map

$$p: \mathbb{R} \rightarrow S^1 \\ t \mapsto (\cos 2\pi t, \sin 2\pi t)$$

the key facts about  $p$  were

path lifting: given  $\gamma: [0,1] \rightarrow S^1$ , then for each  $x \in p^{-1}(\gamma(0))$ ,  $\exists$  unique  $\tilde{\gamma}_x: [0,1] \rightarrow \mathbb{R}$  s.t.  $\tilde{\gamma}_x(0) = x$  and  $p \circ \tilde{\gamma}_x = \gamma$

homotopy lifting: given a homotopy  $H: [0,1] \times [0,1] \rightarrow S^1$  then for each  $x \in p^{-1}(H(0,0))$ ,  $\exists$  unique  $\tilde{H}_x: [0,1] \times [0,1] \rightarrow \mathbb{R}$  s.t.  $\tilde{H}_x(0,0) = x$  and  $p \circ \tilde{H}_x = H$

to prove these properties we used that:

$$S^1 = A \cup B \text{ with } A \text{ and } B \text{ open} \\ p^{-1}(A) = \bigcup_{i=-\infty}^{\infty} A_i, \quad p^{-1}(B) = \bigcup_{i=-\infty}^{\infty} B_i \text{ s.t.}$$

- 1)  $A_i, B_i$  are open in  $\mathbb{R}$
- 2)  $A_i$  are disjoint (same for  $B_i$ 's)
- 3)  $p|_{A_i}: A_i \rightarrow A$  and  $p|_{B_i}: B_i \rightarrow B$  are homeomorphisms

generalizing this we have:

given a topological space  $X$

a covering space of  $X$  is a pair  $(\tilde{X}, p)$  where  $\tilde{X}$  is a

topological space and

$$p: \tilde{X} \rightarrow X$$

is a continuous map (called a covering map) such that

$\forall x \in X$ , there is an open set  $U \subset X$  containing  $x$

$$\text{s.t. } p^{-1}(U) = \{U_\alpha\}_{\alpha \in I}$$

where the  $U_i$  are open, pairwise disjoint sets in  $\tilde{X}$


and  $p|_{U_i}: U_i \rightarrow U$  is a homeomorphism

( $U$  is called an evenly covered set)

examples:


1)  $p: \mathbb{R} \rightarrow S^1$  is clearly a covering map

2)  $p_n: S^1 \rightarrow S^1: \theta \mapsto n\theta$  can easily be seen to be a cover



3)  $p_i: \mathbb{R}^2 \rightarrow T^2 = S^1 \times S^1$  where  $p$  is from 1)  
 $(x, y) \mapsto (p(x), p(y))$

can easily be checked  
to be a covering map



more generally

exercise: if  $p_X: \tilde{X} \rightarrow X$  and  $p_Y: \tilde{Y} \rightarrow Y$  are covering maps,  
then show  $p(x, y) = (p_X(x), p_Y(y))$  is a covering map

$$p: \tilde{X} \times \tilde{Y} \rightarrow X \times Y$$

repeating the proofs we gave for  $p: \mathbb{R} \rightarrow S^1$  (Th<sup>m</sup> IV.8 and 10)

we get

### Th<sup>m</sup> 1 (path lifting):

if  $p: \tilde{X} \rightarrow X$  is a covering map  
 $\gamma: [0,1] \rightarrow X$  is a path and  
 $x \in p^{-1}(\gamma(0))$

then  $\exists$  unique  $\tilde{\gamma}_x: [0,1] \rightarrow \tilde{X}$  such that  
 $\tilde{\gamma}_x(0) = x$  and  $p \circ \tilde{\gamma}_x = \gamma$

### Th<sup>m</sup> 2 (Homotopy lifting):

if  $p: \tilde{X} \rightarrow X$  is a covering map,  
 $H: [0,1] \times [0,1] \rightarrow X$  a homotopy, and  
 $x \in p^{-1}(H(0,0))$

then  $\exists$  unique  $\tilde{H}_x: [0,1] \times [0,1] \rightarrow \tilde{X}$  such that  
 $\tilde{H}_x(0,0) = x$  and  $p \circ \tilde{H}_x = H$

### lemma 3:

let  $p: \tilde{X} \rightarrow X$  be a covering map with  $X$  connected  
if  $\exists$  a point  $x_0 \in X$  with  $|p^{-1}(x_0)| = k$ , then  
 $|p^{-1}(x)| = k, \forall x \in X$

$|p^{-1}(x)|$  is called the degree of the covering space

Proof: let  $A = \{x \in X \text{ s.t. } |p^{-1}(x)| = k\}$

$A \neq \emptyset$  since  $x_0 \in A$

Claim:  $A$  is open

indeed if  $x \in A$ , then let  $U$  be an evenly covered  
open set containing  $x$

$$p^{-1}(U) = \{U_\alpha\}_{\alpha \in I}$$

but  $U_\alpha \cap p^{-1}(x) = 1$  point  $\forall \alpha$

$$\therefore I = \{1, \dots, k\}$$

so  $|p^{-1}(y)| = k, \forall y \in U$

$\therefore A$  is open

Claim:  $A$  is closed

similar argument *exercise*

since  $X$  is connected  $A = X$  (lemma II.10) 

lemma 4:

$p: \tilde{X} \rightarrow X$  a covering map,  $\tilde{x}_0 \in \tilde{X}, x_0 = p(\tilde{x}_0)$   
then

$$p_*: \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$$

is injective

Moreover,  $[\gamma] \in \pi_1(X, x_0)$


$\Leftrightarrow$

the lift of  $\gamma$  to a path based  
at  $\tilde{x}_0$  is a loop in  $\tilde{X}$

Proof:  $[\gamma] \in \pi_1(X, x_0)$

suppose  $p_*([\gamma]) = e$  i.e.  $p \circ \gamma \simeq x_0$

so  $\exists$  homotopy  $H: [0,1] \times [0,1] \rightarrow X$

s.t. 

homotopy lifting says  $\exists \tilde{H}: [0,1] \times [0,1] \rightarrow \tilde{X}$

s.t.  $\tilde{H}(0,0) = \tilde{x}_0$  and  $p \circ \tilde{H} = H$

*constant path* 

note:  $p \circ \tilde{H}(s, 0) = p \circ \gamma$  so  $\tilde{H}(s, 0)$  is a lift of  $p \circ \gamma$  starting at  $\tilde{x}_0$ , so it is  $\gamma$


$$\therefore \tilde{H}(s, 0) = \gamma(s)$$

also  $\tilde{H}(0, t) \in p^{-1}(x_0)$  ← points with discrete topology

$$\text{so } \tilde{H}(0, t) = \tilde{x}_0 \quad \forall t$$

similarly  $\tilde{H}(1, t) = \tilde{x}_0 \quad \forall t$  and  $\tilde{H}(s, 1) = \tilde{x}_0 \quad \forall s$

i.e.



is a homotopy  $\gamma \simeq \tilde{x}_0$

$$\therefore [\gamma] = e$$

and  $p_*$  is injective

now, if  $[\eta] \in p_* (\pi_1(\tilde{X}, \tilde{x}_0))$  then  $\exists [\gamma] \in \pi_1(\tilde{X}, \tilde{x}_0)$

$$\text{s.t. } p_*([\gamma]) = [\eta] \quad \text{i.e. } p \circ \gamma \simeq \eta$$

let  $\tilde{\eta}$  be a lift of  $\eta$  starting at  $\tilde{x}_0$

by homotopy lifting  $\gamma \simeq \tilde{\eta}$  rel end points

but  $\gamma$  a loop so  $\tilde{\eta}$  a loop too

if  $[\eta] \notin p_* (\pi_1(\tilde{X}, \tilde{x}_0))$ , then the lift  $\tilde{\eta}$  of  $\eta$  based at  $\tilde{x}_0$  can't be a loop since if it were then  $[\tilde{\eta}] \in \pi_1(\tilde{X}, \tilde{x}_0)$  and  $[\eta] = p_*([\tilde{\eta}]) \notin \emptyset$

exercise:  $[\pi_1(X, x_0) : p_*(\pi_1(\tilde{X}, \tilde{x}_0))] = \text{degree of } (\tilde{X}, p)$

↑ index of subgroup

Hint: Show there is a bijection from right cosets of  $p_*(\pi_1(\tilde{X}, \tilde{x}_0))$  to  $p^{-1}(x_0)$

examples:

1)  $p: \mathbb{R} \rightarrow S^1$

$$p_*: \pi_1(\mathbb{R}) \rightarrow \pi_1(S^1)$$

$$\begin{matrix} \parallel & \parallel S \\ \{e\} & \mathbb{Z} \end{matrix}$$

no non-trivial loop in  $S^1$  lifts to a loop in  $\mathbb{R}$

degree =  $\infty = [\mathbb{Z} : \{e\}]$

2)  $p_n: S^1 \rightarrow S^1: \theta \mapsto n\theta$


$$(p_n)_*: \pi_1(S^1) \rightarrow \pi_1(S^1)$$

$$\begin{matrix} \parallel S & \parallel S \\ \mathbb{Z} & \mathbb{Z} \\ \downarrow & \downarrow \\ m & \rightarrow nm \end{matrix}$$

so  $\text{im}(p_n)_* = n\mathbb{Z}$

degree =  $n = [\mathbb{Z} : n\mathbb{Z}]$

loop in  $S^1$  lifts to a loop iff it "goes around"  $S^1$  a multiple of  $n$  times



3)  $X =$

$$\tilde{X} =$$

$$\xrightarrow{p} \text{ } = X$$


exercise: this is a covering map

note:  $\pi_1(\tilde{X}) = F_3$  generated by  $\tilde{a}_1, \tilde{b}_1 * \tilde{b}_2, \tilde{b}_1 * \tilde{a}_2 * \tilde{b}_1$

so  $\text{image}(p_*) = \langle a, b^2, bab^{-1} \rangle = G$


$G$  has index 2 in  $\pi_1(X, x_0) \cong F_2$ !

note rank went up!

4) consider  $\Sigma_2$  

let's find a degree 2 cover (there are actually a lot)

consider



define  $p$  to be

exercise:

1) show this is a 2-fold covering map  $\Sigma_3 \rightarrow \Sigma_2$

2) work out  $\text{im}(p_*)$

3) Experiment constructing other covers of other surfaces


e.g.  $\Sigma_n \rightarrow \Sigma_2$  by an  $n-1$  fold cover for  $n \geq 2$

let  $p: \tilde{X} \rightarrow X$  be a covering map with  $p(\tilde{x}_0) = x_0$

$f: Y \rightarrow X$  be a continuous map such that  $f(y_0) = x_0$

a lift of  $f$  to  $\tilde{X}$  is a continuous map  $\tilde{f}: Y \rightarrow \tilde{X}$

s.t.  $\tilde{f}(y_0) = \tilde{x}_0$  and  $p \circ \tilde{f} = f$



Th<sup>m</sup> 5 (lifting criterion):

$p: \tilde{X} \rightarrow X$  a covering map,  $p(\tilde{x}_0) = x_0$

$f: Y \rightarrow X$  a continuous map s.t.  $f(y_0) = x_0$

assume  $Y$  is path connected and

locally path connected

Then  $\exists$  a lift  $\tilde{f}: Y \rightarrow \tilde{X}$  of  $f$


$\Leftrightarrow$

$$f_*(\pi_1(Y, y_0)) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$$

if  $\tilde{f}$  exists it is unique

a space is locally path connected if for every point  $x$  and open set  $U$  containing it, there is an open set  $V$  such that  $x \in V \subset U$  and  $V$  is path connected

example:



$$(\{y_n\} \times [0, 1]) \cup (\{0\} \times [0, 1]) \cup ([0, 1] \times \{0\})$$

path connected but not  
locally path connected

note: all manifolds are locally path connected



Proof:  $(\Rightarrow)$  if  $\tilde{f}$  exists, then clearly

$$f_* (\pi_1(Y, y_0)) = p_* \circ \tilde{f}_* (\pi_1(Y, y_0)) \subseteq p_* (\pi_1(\tilde{X}, \tilde{x}_0))$$

$(\Leftarrow)$  need to construct  $\tilde{f}: Y \rightarrow \tilde{X}$

given  $y \in Y$ , let  $\gamma_y: [0, 1] \rightarrow Y$  be a path s.t.

$$\gamma_y(0) = y_0, \gamma_y(1) = y \quad (\text{use path connected})$$

$f \circ \gamma_y$  is a path in  $X$  from  $x_0 = f(y_0)$  to  $f(y)$

lift  $f \circ \gamma_y$  to a path  $\tilde{\gamma}_y$  in  $\tilde{X}$  starting at  $\tilde{x}_0$

$$\text{define: } \tilde{f}(y) = \tilde{\gamma}_y(1)$$


if  $\tilde{f}$  is well-defined, then clearly  $p \circ \tilde{f}(y) = f(y)$

so  $\tilde{f}$  is a lift of  $f$

to see  $\tilde{f}$  is well-defined, let  $\gamma'_y$  be another path from  $y_0$  to  $y$

note:  $\gamma_y * \overline{\gamma'_y}$  is a loop in  $Y$  based at  $y_0$

$$\text{so } [\gamma_y * \overline{\gamma'_y}] \in \pi_1(Y, y_0)$$



$$\text{and } f_* [\gamma_y * \overline{\gamma'_y}] = [(f \circ \gamma_y) * \overline{(f \circ \gamma'_y)}] \in \pi_1(X, x_0)$$

$$\text{by assumption } [(f \circ \gamma_y) * \overline{(f \circ \gamma'_y)}] \in p_* (\pi_1(\tilde{X}, \tilde{x}_0))$$

so by lemma 4  $(f \circ \gamma_y) * \overline{(f \circ \gamma'_y)}$  lifts to a loop


$$\text{in } \tilde{X} \text{ based at } x_0: \overline{(f \circ \gamma_y) * \overline{(f \circ \gamma'_y)}}$$

can easily check by uniqueness of lifts that

$$\text{this loop is } \overline{(f \circ \gamma_y) * \overline{(f \circ \gamma'_y)}} \quad \text{lift starting at } \tilde{\gamma}_y(1)$$

so  $\tilde{f}$  is well-defined  
 the last thing we need to do is see  $\tilde{f}$  is continuous.  
 this is more involved (and uses local connectivity)  
 you can find a proof in Hatcher, but the idea is:

given  $y \in Y$ ,  $\exists$  an open set  $U \subset Y$  containing  
 $y$  and open set  $V$  in  $\tilde{X}$  containing  
 $\tilde{f}(y)$  such that  $\tilde{f}|_U = \underbrace{p|_V^{-1}}_{\text{continuous}} \circ f$



Fact: given a surface  $\Sigma_g$  of genus  $g$

if  $g > 0$ , then  $\exists$  a covering map

$$p: \mathbb{R}^2 \rightarrow \Sigma_g$$

(for  $g > 1$ , this uses "hyperbolic geometry")

Thm 6:

If  $g \geq 1$  and  $n \geq 2$ , then any

$$f: S^n \rightarrow \Sigma_g$$

is homotopic to the constant map!

Recall, this was used in the proof of Thm VI.6

Proof: given  $f$ , clearly  $f_*(\pi_1(S^n)) = \{e\} \subset p_*(\pi_1(\mathbb{R}^2))$

so  $f$  lifts to a map  $\tilde{f}: S^n \rightarrow \mathbb{R}^2$   
 by Thm 5

↑ covering map  
 above

$$\text{let } \tilde{H}: S^n \times [0,1] \rightarrow \mathbb{R}^2$$

$$(p,t) \mapsto t\tilde{f}(p)$$

$$\tilde{H}(p,0) = \text{constant}$$

$$\tilde{H}(p,1) = \tilde{f}$$

$$\text{set } H = p_0 \tilde{H}: [0,1] \times [0,1] \rightarrow \Sigma_g$$

this is a homotopy from the constant map to  $f$  

We saw that for every covering  $p: \tilde{X} \rightarrow X$ , there is a subgroup  $G = p_*(\pi_1(\tilde{X}, \tilde{x}_0))$  of  $\pi_1(X, x_0)$

For most spaces, there is a converse!

Fact:

let  $X$  be path connected

locally path connected

semi-locally simply connected

Then  $\forall G < \pi_1(X, x_0)$  there is a covering space

$$p: \tilde{X} \rightarrow X \text{ such that } p_*(\pi_1(\tilde{X}, \tilde{x}_0)) = G$$

a space  $X$  is semi-locally simply connected if

$\forall x \in X, \exists$  an open set  $U \subset X$  such that  $x \in U$  and

$$\iota_*: \pi_1(U, x) \rightarrow \pi_1(X, x_0)$$

is the trivial map, where  $\iota: U \rightarrow X$  is inclusion

Fact: manifolds and CW complexes are semi-locally simply connected.

example:



is not

We will not prove this, but the idea for  $G = \{e\} < \pi_1(X)$  is

let  $\tilde{X} = \{\text{paths in } X \text{ starting at } x_0\} / \sim$

here  $\gamma \sim \eta$  if they are homotopic  
rel end points

set  $p: \tilde{X} \rightarrow X: [\gamma] \mapsto \gamma(1)$

you can put a topology on  $\tilde{X}$  so this is  
the desired covering space

## B. Subgroups

we use covering spaces to show

Th<sup>m</sup> 7 (Nielsen-Schreier):

any subgroup of a free group is free

We need some lemmas

lemma 8:


let  $X$  be a graph, then  $\pi_1(X)$  is free

Proof: we can assume  $X$  is connected

if  $X$  has only one vertex, then  $X$  is a wedge of circles

so from Section VI we know  $\pi_1(X)$  free group

if  $X$  has more than one vertex, then there is an  
edge  $e$  in  $X$  connecting distinct vertices



CW Fact: if  $X$  is a CW complex, and  $A$  is a contractible subcomplex, then  $X/A \cong X$

exercise: try to prove this in above situation

so  $X/e \cong X$  and  $\pi_1(X) \cong \pi_1(X/e)$ ,

but  $X/e$  is a graph with one less vertex

thus we can inductively find a graph  $Y$  with one vertex that is homotopy equivalent to  $X$

$\therefore$  done 

lemma 9:

If  $X$  is a graph and  $p: \tilde{X} \rightarrow X$  is a covering space then  $\tilde{X}$  is a graph

more generally, coverings of CW complexes are CW complexes

Sketch of Proof:

$p^{-1}(X^{(0)})$  is a discrete set of points in  $\tilde{X}$


$\leftarrow$  0-skeleton  $\nearrow$  this will be  $\tilde{X}^{(0)}$

each edge  $e$  of  $X$  is a path so it lifts to  $\tilde{X}$

the union of all lifts of all edges will be the edges of  $\tilde{X}$

to make this rigorous we need to see how to

"attach" edges to the vertices

but hopefully this is intuitively clear 

## Proof of Th<sup>m</sup> 7:

given a free group  $F_n$  on  $n$  generators

let  $W_n =$  wedge of  $n$ -circles

$$\text{so } \pi_1(W_n) \cong F_n$$

given any  $G < F_n \cong \pi_1(W_n)$ ,  $\exists$  a covering space

$$p: \tilde{X} \rightarrow X$$

such that  $\pi_1(\tilde{X}) \cong p_*(\pi_1(\tilde{X})) = G$  ↙ by injectivity

now lemma 9 says  $\tilde{X}$  is a graph

and thus by lemma 8,  $\pi_1(\tilde{X})$  is a free group

$\therefore G$  is a free group 