

# E Covering Spaces

a covering space of a space  $X$  is a pair  $(\tilde{X}, p)$  where

$\tilde{X}$  is a space and

$p: \tilde{X} \rightarrow X$  is a continuous map such that every point  $x \in X$  has an evenly covered neighborhood

an open set  $U$  is called evenly covered if

$p^{-1}(U) =$  disjoint union of open sets  $\{U_\alpha\}$  in  $\tilde{X}$

such that  $p|_{U_\alpha}: U_\alpha \rightarrow U$  is a homeomorphism  $\forall \alpha$

## examples:

1) a homeomorphism  $p: \tilde{X} \rightarrow X$  is a covering space


2)  $p: \mathbb{R} \rightarrow S^1: t \mapsto (\cos 2\pi t, \sin 2\pi t)$  is a covering space

since we showed  $A = S^1 - \{(1,0)\}$  and  $B = S^1 - \{(-1,0)\}$  are evenly covered


exercise: if  $(\tilde{X}, p)$  is a covering space of  $X$  and  $(\tilde{Y}, q)$  " " " "  $Y$


then  $(\tilde{X} \times \tilde{Y}, p \times q)$  is a covering space of  $X \times Y$

so  $p: \mathbb{R}^2 \rightarrow T^2: (x, y) \mapsto ((\cos 2\pi x, \sin 2\pi x), (\cos 2\pi y, \sin 2\pi y))$  is a covering space of  $T^2$




3)  $p_n: S^1 \rightarrow S^1: \theta \mapsto n\theta$





4)  $X =$   wedge of 2 circles




note: a evenly covered  
b evenly covered

$x_0$  has nbhd  $X_{x_0}$


evenly covered by 

exercise: explicitly write out  $p$

similarly another cover of  $X$  is



5)  $\mathbb{R}P^2 = S^2 / \sim$  points in  $S^2 \sim$  to antipode



similarly  $S^n \rightarrow \mathbb{R}P^n$  a covering map

### lemma 19:

let  $(\tilde{X}, p)$  be a covering space of a connected space  $X$   
 the cardinality  $|p^{-1}(x)|$  is independent of  $x \in X$

$|p^{-1}(x)|$  is called the degree of the covering space

Proof: for some  $x_0 \in X$ , let  $k = |p^{-1}(x_0)|$

let  $A = \{x \in X : |p^{-1}(x)| = k\}$  note:  $A \neq \emptyset$  since  $x_0 \in A$

if  $x \in A$  then let  $U$  be a evenly covered nbhd of  $x$

$$\text{so } p^{-1}(U) = \{U_1, \dots, U_k\}$$

$$\therefore |p^{-1}(x')| = k \quad \forall x' \in U$$

$$\therefore U \subset A \text{ and } A \text{ open}$$


similarly  $X - A$  open so  $A$  closed

$\therefore X = A$  since  $X$  connected  $\square$

If  $(\tilde{X}, p)$  a covering space of  $X$

$f: Y \rightarrow X$  a continuous map

then  $\tilde{f}: Y \rightarrow \tilde{X}$  a lift of  $f$  if  $p \circ \tilde{f} = f$



if  $f(y_0) = x_0$  and  $\tilde{x}_0 \in \tilde{X}$  s.t.  $p(\tilde{x}_0) = x_0$ , then  $\tilde{f}$  is a lift of  $f$  based at  $\tilde{x}_0$   
 if it is a lift and  $\tilde{f}(y_0) = \tilde{x}_0$

### lemma 20:

$(\tilde{X}, p)$  a covering space of  $X$ ,  $x_0 \in X$  and  $\tilde{x}_0 \in p^{-1}(x_0)$

path lifting  $\rightarrow$

a) each path  $\gamma: [0, 1] \rightarrow X$  based at  $x_0$  has a unique lift to a path  $\tilde{\gamma}: [0, 1] \rightarrow \tilde{X}$  based at  $\tilde{x}_0$

homotopy lifting  $\rightarrow$

b) if  $H: Y \times [0, 1] \rightarrow X$  is a homotopy with  $h_0(y) = H(y, 0)$  and  $\tilde{h}_0: Y \rightarrow \tilde{X}$  a lift of  $h_0$  then  $\exists$  a unique homotopy

$$\tilde{H}: Y \times [0, 1] \rightarrow \tilde{X} \text{ s.t. } \tilde{h}_0(y) = \tilde{H}(y, 0)$$

Proof: same as proof of lemma 12 for a) and b) if  $Y = [0, 1]$   
 for general case see Th<sup>m</sup> 23 below (exercise)  $\square$

lemma 21:

If  $(\tilde{X}, p)$  is a path connected covering space of  $X$   
and  $x_0 \in X, \tilde{x}_0 \in p^{-1}(x_0)$

then  $p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$

(1) is injective

(2) its image is the set of loops in  $\pi_1(X, x_0)$  that when lifted to paths in  $\tilde{X}$  based at  $\tilde{x}_0$ , they are loops

(3)  $[\pi_1(X, x_0) : \pi_1(\tilde{X}, \tilde{x}_0)] = \text{degree of } (\tilde{X}, p)$  if  $\tilde{X}$  is path connected

examples:

1)  $p: \mathbb{R} \rightarrow S^1$

$$p_* : \pi_1(\mathbb{R}) \rightarrow \pi_1(S^1)$$

$$\begin{matrix} \text{"} & \text{S11} \\ \{e\} & \mathbb{Z} \end{matrix}$$

no non-trivial loop in  $S^1$  lifts to a loop in  $\mathbb{R}$   
degree =  $\infty = [\mathbb{Z} : \{0\}]$


2)  $p_n: S^1 \rightarrow S^1$

$$(p_n)_* : \pi_1(S^1) \rightarrow \pi_1(S^1)$$

$$\begin{matrix} \text{115} & \text{115} \\ \mathbb{Z} & \rightarrow \mathbb{Z} \\ 1 & \longmapsto n \end{matrix}$$



so  $\text{im}(p_n)_* = n\mathbb{Z}$

degree  $p_n = n = [\mathbb{Z} : n\mathbb{Z}]$



any loop in  $\uparrow$  that wraps a multiple of 3 times lifts to a loop

Proof: 1) Suppose  $p_*([\gamma]) = [e]$



note:  $\tilde{H}(0, t)$  lift of  $H(0, t) = x_0$  so constantly  $\tilde{x}_0$

why is this  $\gamma$ ?

similarly  $\tilde{H}(1,t) = \tilde{x}_0 = \tilde{H}(s,1)$

so  $\gamma \sim e_{\tilde{x}_0}$  by  $\tilde{H}$  and  $[\gamma] = [e]$

(2) clearly if  $[\gamma] \in \pi_1(X, x_0)$  and  $\gamma$  lifts to a loop  $\tilde{\gamma}$  based at  $\tilde{x}_0$   
then  $p_*([\tilde{\gamma}]) = \gamma$

and if  $[\gamma] = p_*([\eta])$ , then  $\gamma \sim p \circ \eta$   $\therefore$  by lemma 20.6)

the lift  $\tilde{\gamma}$  of  $\gamma$  based at  $\tilde{x}_0$  is homotopic to  $\eta$   
(rel end pts)

so  $\tilde{\gamma}$  a loop

(3) let  $H = p_* (\pi_1(\tilde{X}, \tilde{x}_0))$

if  $[\gamma] \in \pi_1(X, x_0)$  and  $[S] \in H$ , then note

if  $\tilde{\gamma}$  a lift of  $\gamma$  based at  $\tilde{x}_0$

$\tilde{\delta} * \tilde{\gamma}$  " "  $\delta * \gamma$  " "  $\tilde{x}_0$

then  $\tilde{\gamma}(1) = \tilde{\delta} * \tilde{\gamma}(1)$  since  $\tilde{\delta}$  is a loop (and  $\tilde{\delta} * \tilde{\gamma} = \tilde{\delta} * \tilde{\gamma}$ )

$\therefore$  we get a map

$$\begin{aligned} \phi: \{ \text{right cosets of } H \} &\longrightarrow p^{-1}(x_0) \\ H[\gamma] &\longmapsto \tilde{\gamma}(1) \end{aligned}$$

that is well-defined

if  $\tilde{x}_1 \in p^{-1}(x_0)$  then let  $\tilde{\delta}$  be a path  $\tilde{x}_0$  to  $\tilde{x}_1$

$\gamma = p \circ \tilde{\delta}$  is a loop in  $X$  based at  $x_0$

and  $\phi(H[\gamma]) = \tilde{\gamma}(1) = \tilde{x}_1$

so  $\phi$  onto

now suppose  $\phi(H[\gamma]) = \phi(H[\eta])$


so if  $\tilde{\gamma}, \tilde{\eta}$  are lifts of  $\gamma, \eta$  based at  $\tilde{x}_0$

then  $\tilde{\gamma}(1) = \tilde{\eta}(1)$

$\therefore \tilde{\gamma} * \tilde{\eta}$  is a loop in  $\tilde{X}$  and so  $p_*([\tilde{\gamma} * \tilde{\eta}]) \in H$

but  $p_*([\tilde{\gamma} * \tilde{\eta}]) = [\gamma] * [\eta]^{-1} \Rightarrow H[\gamma] = H[\eta]$

i.e.  $\phi$  is one-to-one



lemma 22:

If  $(\tilde{X}, p)$  is a path connected covering space of  $X$  and  $x_0 \in X$   
 then  $\{p_*(\pi_1(\tilde{X}, \tilde{x}_i))\}_{\tilde{x}_i \in p^{-1}(x_0)}$   
 is a conjugacy class of subgroups of  $\pi_1(X, x_0)$

Proof: let  $\tilde{x}_0, \tilde{x}_i \in p^{-1}(x_0)$  and set  $H_i = p_*(\pi_1(\tilde{X}, \tilde{x}_i)) \quad i=0,1$

let  $h: [0,1] \rightarrow \tilde{X}$  be a path  $\tilde{x}_0$  to  $\tilde{x}_i$

then  $\gamma = p \circ h$  is a loop in  $X$

if  $[\eta] \in H_i$  then  $\eta$  lifts to a loop  $\tilde{\eta}$

based at  $\tilde{x}_i$  (by lemma 21)

so  $h * \tilde{\eta} * \bar{h}$  is a loop based at  $\tilde{x}_0$

$$\begin{aligned} \therefore [\gamma] \cdot [\eta] \cdot [\gamma]^{-1} &= [(p \circ h) * (p \circ \tilde{\eta}) * \overline{(p \circ h)}] \\ &= p_* [h * \tilde{\eta} * \bar{h}] \in H_0 \end{aligned}$$

$$\therefore [\gamma] H_i [\gamma]^{-1} \subseteq H_0$$

similarly  $[\gamma]^{-1} H_0 [\gamma] \subseteq H_i \quad \therefore H_0 = [\gamma] H_i [\gamma]^{-1}$

now if  $H$  is conjugate to  $H_0$ , then  $\exists [\alpha] \in \pi_1(X, x_0)$  s.t.

$$[\alpha] H [\alpha]^{-1} = H_0$$


if  $[\alpha] \in H_0$  then  $H = H_0$  and done

if  $[\alpha] \notin H_0$  then  $\alpha$  lifts to a path  $\tilde{\alpha}$  starting at  $\tilde{x}_0$

let  $\tilde{x}_i = \tilde{\alpha}(1)$

set  $H_i = p_*(\pi_1(\tilde{X}, \tilde{x}_i))$

from above  $H_0 = [\alpha] H_i [\alpha]^{-1} \quad \therefore H = H_i$



Thm 23:

let  $(\tilde{X}, p)$  be a covering space of  $X$ ,  $\tilde{x}_0 \in \tilde{X}$  and  $x_0 = p(\tilde{x}_0)$

suppose  $f: Y \rightarrow X$  is a continuous map with

$Y$  path connected and locally path connected

$$y_0 \in Y \text{ st. } f(y_0) = x_0$$

Then  $\exists$  a lift  $\tilde{f}: Y \rightarrow \tilde{X}$  ( $\rho \circ \tilde{f} = f$ ) st.  $\tilde{f}(y_0) = \tilde{x}_0$


$\Leftrightarrow$

$$f_* (\pi_1(Y, y_0)) \subseteq p_* (\pi_1(\tilde{X}, \tilde{x}_0))$$

(and if lift exists it is unique, see lemma 24 below)

$Y$  is locally path connected if  $\forall y \in Y$  and open sets  $U$  containing  $y$ ,  $\exists$  an open set  $V$  st.  $y \in V \subset U$  and  $V$  is path connected

example: the comb space  $C = (\{\frac{1}{n}\} \times [0,1]) \cup \{0\} \times [0,1] \cup [0,1] \times \{0\}$  is path connected but not locally path connected



Proof:  $(\Rightarrow) f_* (\pi_1(Y, y_0)) = p_* (\tilde{f}_* (\pi_1(Y, y_0))) \subseteq p_* (\pi_1(\tilde{X}, \tilde{x}_0))$

$(\Leftarrow)$  for  $y \in Y$  let  $\gamma$  be a path  $y_0$  to  $y$

so  $f \circ \gamma$  is a path in  $X$  based at  $x_0$

$\exists!$  lift  $\tilde{f} \circ \gamma: [0,1] \rightarrow \tilde{X}$  based at  $\tilde{x}_0$

define:  $\tilde{f}(y) = \tilde{f} \circ \gamma(1)$  note: if well-defined, then clearly  $\rho \circ \tilde{f} = f$  and  $\tilde{f}(y_0) = \tilde{x}_0$

Claim:  $\tilde{f}$  well-defined


let  $\gamma, \eta$  be two paths  $y_0$  to  $y$

$\gamma * \bar{\eta}$  is a loop in  $Y$  based at  $y_0$

so  $f_* ([\gamma * \bar{\eta}]) \in p_* (\pi_1(\tilde{X}, \tilde{x}_0))$

i.e.  $(f \circ \gamma) * (f \circ \bar{\eta})$  lifts to a loop in  $\tilde{X}$  based at  $\tilde{x}_0$

but  $\widetilde{(f \circ \gamma) * (f \circ \bar{\eta})} = \underbrace{\tilde{f} \circ \gamma}_{\text{lift based at } \tilde{x}_0} * \underbrace{\tilde{f} \circ \bar{\eta}}_{\text{lift based at } \tilde{f} \circ \gamma(1)}$




note: 1)  $\tilde{f} \circ \bar{\eta}(1) = \tilde{x}_0$

2)  $\widetilde{f \circ \eta} = \widetilde{f \circ \gamma}$  so  $\tilde{f} \circ \eta(1) = \tilde{f} \circ \gamma(1)$  and  $\tilde{f}$  is well-defined!

Claim:  $\tilde{f}$  is continuous

given an open set  $U \subset \tilde{X}$  we show  $\forall y \in \tilde{f}^{-1}(U), \exists$  an open set  $V$   
 st.  $y \in V \subset \tilde{f}^{-1}(U)$   
 $(\therefore \tilde{f}^{-1}(U) \text{ open})$



idea is on small  
 nbhd of  $y$ ,  $\tilde{f}$  is  
 $f \circ (p|_W)^{-1}$

let  $W$  be an evenly covered nbhd of  $f(y)$

and  $\tilde{W}$  open set in  $\tilde{X}$  s.t.  $\tilde{f}(y) \subset \tilde{W} \subset U$

and  $p|_{\tilde{W}}: \tilde{W} \rightarrow W$  homeomorphism (might need to shrink  $W$ )

$Y$  locally path connected  $\Rightarrow \exists V$  open in  $Y$  s.t.  $y \in V \subset \tilde{f}^{-1}(W)$   
 and  $V$  path connected

now fix a path  $\delta$  from  $y_0$  to  $y$

for any  $y' \in V$  let  $\eta$  be a path  $y$  to  $y'$  in  $V$

so  $\delta * \eta$  is a path  $y_0$  to  $y'$

$$\therefore \tilde{f}(y') = \widetilde{f \circ (\delta * \eta)}(1)$$

but if  $\tilde{f} \circ \eta$  is lift of  $f \circ \eta$  based at  $\tilde{f} \circ \delta(1) = \tilde{f}(y)$

$$\text{then } \widetilde{f \circ (\delta * \eta)}(1) = \tilde{f} \circ \eta(1)$$

$$\text{and we know } \tilde{f} \circ \eta = \underbrace{(p|_W)^{-1}}_{\leftarrow \text{since this is a lift and lift is unique!}} \circ f \circ \eta$$

$$\therefore \tilde{f}(y') \in \tilde{W} \subset U$$

$$\text{i.e. } V \subset \tilde{f}^{-1}(U) \quad \square$$

### lemma 24:

$(\tilde{X}, p)$  a covering space of  $X$

let  $\tilde{f}_1, \tilde{f}_2: Y \rightarrow \tilde{X}$  be two lifts of  $f: Y \rightarrow X$

if  $Y$  is connected and  $\tilde{f}_1$  and  $\tilde{f}_2$  agree at one point then  $\tilde{f}_1 = \tilde{f}_2$



Proof:

$$\text{let } A = \{y \in Y \text{ s.t. } \tilde{f}_1(y) = \tilde{f}_2(y)\}$$

$A \neq \emptyset$  by assumption

if  $y \in A$  then let  $U$  be an evenly covered nbhd of  $f(y)$

$$\text{let } \tilde{U} \text{ be an open set in } \tilde{X} \text{ s.t. } \tilde{f}_1(y) = \tilde{f}_2(y) \in \tilde{U}$$

and  $p|_{\tilde{U}}: \tilde{U} \rightarrow U$  is a homeomorphism

since  $f$  is continuous  $\exists$  open nbhd  $V$  of  $y$  s.t.  $f(V) \subset U$

$$\text{now } \tilde{f}_1|_V = (p|_{\tilde{U}})^{-1} \circ f|_V = \tilde{f}_2|_V$$

$\therefore V \subset A$  and  $A$  open

if  $y \notin A$ , then with  $U$  as above,  $\exists \tilde{U}_1, \tilde{U}_2$  open in  $\tilde{X}$  s.t.

$$\tilde{f}_1(y) \in \tilde{U}_1 \text{ and } p|_{\tilde{U}_1}: \tilde{U}_1 \rightarrow U \text{ a homeomorphism}$$

$$\text{clearly } \tilde{U}_1 \cap \tilde{U}_2 = \emptyset$$

$$\therefore \text{if } V \text{ is as above, then } \tilde{f}_1(V) \subset \tilde{U}_1$$

so  $X - A$  open

$\therefore$  by connectedness of  $X$ ,  $A = X$   $\square$

Two covering spaces  $(\tilde{X}_1, p_1), (\tilde{X}_2, p_2)$  of  $X$  are isomorphic if  $\exists$  a

homeomorphism  $h: \tilde{X}_1 \rightarrow \tilde{X}_2$  s.t.  $p_2 \circ h = p_1$

$$\begin{array}{ccc} \tilde{X}_1 & \xrightarrow{h} & \tilde{X}_2 \\ p_1 \downarrow & & \downarrow p_2 \\ & X & \end{array}$$

note: this is an equivalence relation

Cor 25:

Suppose  $(\tilde{X}_1, p_1), (\tilde{X}_2, p_2)$  are path connected, locally path connected covering spaces of  $X$ ,  $x_0 \in X$ ,  $\tilde{x}_i \in p_i^{-1}(x_0)$

$$\text{a) if } (p_1)_*(\pi_1(\tilde{X}_1, \tilde{x}_1)) \subset (p_2)_*(\pi_1(\tilde{X}_2, \tilde{x}_2))$$

then  $p_1$  lifts to a covering space  $p: \tilde{X}_1 \rightarrow \tilde{X}_2$  taking  $\tilde{x}_1$  to  $\tilde{x}_2$ .


b)  $(\tilde{X}_1, \tilde{x}_1)$  and  $(\tilde{X}_2, \tilde{x}_2)$  are isomorphic covering spaces of  $X$  by an isomorphism taking  $\tilde{x}_1$  to  $\tilde{x}_2$

$\Leftrightarrow$

$$(p_1)_*(\pi_1(\tilde{X}_1, \tilde{x}_1)) = (p_2)_*(\pi_1(\tilde{X}_2, \tilde{x}_2))$$

c)  $(\tilde{X}_1, p_1)$  and  $(\tilde{X}_2, p_2)$  are isomorphic covering spaces of  $X$   
 $\Leftrightarrow$   
 $(p_1)_*(\pi_1(\tilde{X}_1, \tilde{x}_1))$  is conjugate to  $(p_2)_*(\pi_1(\tilde{X}_2, \tilde{x}_2))$

Proof: a) by Th<sup>m</sup> 23 we get a lift  $p: \tilde{X}_1 \rightarrow \tilde{X}_2$  taking  $\tilde{x}_1$  to  $\tilde{x}_2$



we now show  $p$  a covering map

$x \in \tilde{X}_2$  let  $U$  be a nbhd of  $p_2(x)$  in  $X$  that is evenly covered by  $p_1$  and  $p_2$

so  $\exists$  a unique  $\tilde{U}$  in  $\tilde{X}_2$  st.  $x \in \tilde{U}$  and

$p_2|_{\tilde{U}}: \tilde{U} \rightarrow U$  is a homeomorphism

let  $p^{-1}(U) = \bigcup_{\alpha} \tilde{U}_{\alpha}$ , clearly  $\bigcup_{\alpha} \tilde{U}_{\alpha} \subset \tilde{p}_1^{-1}(U)$  so  $p_1|_{\tilde{U}_{\alpha}}: \tilde{U}_{\alpha} \rightarrow U$  a homeo.


so  $p|_{\tilde{U}_{\alpha}} = p_2^{-1}|_{U_{\alpha}} \circ p_1|_{\tilde{U}_{\alpha}}$  is a homeomorphism  $\tilde{U}_{\alpha} \rightarrow \tilde{U}$

$\therefore$  each point in  $p$  has an evenly covered nbhd. ✓


b)  $(\Rightarrow)$  clear

$(\Leftarrow)$  let  $\tilde{p}_1: \tilde{X}_1 \rightarrow \tilde{X}_2$  be lift of  $p_1$

$\tilde{p}_2: \tilde{X}_2 \rightarrow \tilde{X}_1$  be lift of  $p_2$



note:  $\tilde{p}_1 \circ \tilde{p}_2: \tilde{X}_2 \rightarrow \tilde{X}_2$  takes  $\tilde{x}_2$  to  $\tilde{x}_2$  and is a lift of  $p_2$  to  $\tilde{X}_2$



but so is  $id_{\tilde{X}_2}: \tilde{X}_2 \rightarrow \tilde{X}_2 \therefore$  by lemma 24,  $\tilde{p}_1 \circ \tilde{p}_2 = id_{\tilde{X}_2}$

similarly  $\tilde{p}_2 \circ \tilde{p}_1 = id_{\tilde{X}_1}$

$\therefore \tilde{p}_1$  and  $\tilde{p}_2$  are homeomorphisms. ✓

c) clear from lemma 22 and b) 

A space  $X$  is semilocally simply connected if each point  $x \in X$  has a neighborhood  $U$  s.t.  $\pi_1(U, x) \rightarrow \pi_1(X, x)$  is trivial  
 ↑  
 induced by inclusion

example:

$$X = \bigcup_{n=1}^{\infty} S_{\frac{1}{n}}(\frac{1}{n}, 0)$$

← circle of radius  $\frac{1}{n}$  about  $(\frac{1}{n}, 0)$

is not semi-locally simply connected

but CW-complexes and manifolds are

note: the cone  $CX$  is not locally simply connected but it is semilocally simply connected

Th<sup>m</sup> 26:

let  $X$  be a path connected,  
 locally path connected, and  
 semilocally simply connected space

$x_0 \in X$ , Then there is a one-to-one correspondence

$$\left\{ \begin{array}{l} \text{base point preserving isomorphism} \\ \text{classes of coverings } (\tilde{X}, p, \tilde{x}_0) \text{ of } (X, x_0) \end{array} \right\} \longleftrightarrow \left\{ \text{subgroups of } \pi_1(X, x_0) \right\}$$

$$\begin{array}{ccc} (\tilde{X}, p, \tilde{x}_0) & \xrightarrow{\quad} & p_* (\pi_1(\tilde{X}, \tilde{x}_0)) \cong \pi_1(\tilde{X}, \tilde{x}_0) \\ (\tilde{X}_H, p_H, \tilde{x}_H) & \xleftarrow{\quad} & H \end{array}$$

- such that
- 1) if  $H < K$ , then  $(\tilde{X}_H, p_H, \tilde{x}_H)$  is also a cover of  $(\tilde{X}_K, p_K, \tilde{x}_K)$
  - 2) if  $p_i$  in  $(\tilde{X}_i, p_i, \tilde{x}_i)$  lifts to a cover of  $(\tilde{X}_2, p_2, \tilde{x}_2)$  taking  $\tilde{x}_i$  to  $\tilde{x}_2$  then  $(p_i)_* (\pi_1(\tilde{X}_i, \tilde{x}_i)) < (p_2)_* (\pi_1(\tilde{X}_2, \tilde{x}_2))$
  - 3)  $[\pi_1(X, x_0) : H] = n \iff (\tilde{X}_H, p_H, \tilde{x}_H)$  is a cover of degree  $n$


in addition, we have the one-to-one correspondence

$$\left\{ \begin{array}{l} \text{isomorphism classes of} \\ \text{coverings } (\tilde{X}, p) \text{ of } X \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{conjugacy classes of} \\ \text{subgroups of } \pi_1(X, x_0) \end{array} \right\}$$

**This is an amazing th<sup>m</sup>!** There is a "lattice" of subgroups of  $\pi_1(X, x_0)$  and a "lattice" of covering spaces of  $X$ . These lattices are the same!  
 we will see there is more to this correspondence later

The unique simply connected cover of  $X$  is called the universal cover

example:



Proof:

note 1) and 2) follow from Cor 25 once we have the one-to-one corresp.  
3) follows from lemma 21

also, once we have a well-defined map  $H < \pi_1(X, x_0) \mapsto (X_H, p_H, \tilde{x}_H)$

$$\text{such that } p_*[\pi_1(\tilde{X}_H, \tilde{x}_H)] = H$$


the fact that the 1<sup>st</sup> correspondance is one-to-one follows from Cor 25, b)  
and the 2<sup>nd</sup> correspondance from Cor 25, c)

so we are left to construct  $(X_H, p_H, \tilde{x}_H)$  given  $H < \pi_1(X, x_0)$

to this end we call two paths  $\gamma, \eta: [0, 1] \rightarrow X$  based at  $x_0$ , H-equivalent

if 1)  $\gamma(1) = \eta(1)$  and

$$2) [\gamma * \bar{\eta}] \in H$$



exercise: 1) This is an equivalence relation

2) if  $H = \{e\}$  then this is just homotopy rel end points

let  $\langle \gamma \rangle$  denote the equivalence class of  $\gamma$

Set  $\tilde{X}_H = \{ \langle \gamma \rangle \mid \gamma \text{ a path in } X \text{ based at } x_0 \}$

(this is just a set, but we put a topology on it later)

$$p_H: \tilde{X}_H \rightarrow X: \langle \gamma \rangle \mapsto \gamma(1)$$

$$\tilde{x}_H = \langle e_{x_0} \rangle$$

note:  $p_H$  is onto since any point  $x \in X$  is connected to  $x_0$  by a path

We want to define a topology on  $\tilde{X}_H$ , but first we need to understand something about the topology on  $X$

Claim:  $\mathcal{U} = \{ U \subset X : U \text{ open, path-connected and } \pi_1(U, x) \rightarrow \pi_1(X, x) \text{ trivial, some } x \in U \}$   
 is a basis for the topology on  $X$

(recall, a collection of open sets in  $X$  is a basis  
 for the topology on  $X$  if  $\forall x \in X$  and open set  $U$  with  $x \in U$   
 $\exists$  an open set  $O$  in the collection st.  $x \in O \subset U$ .  
 i.e. any open set is a union of sets in the collection)

Pf: note: if  $\pi_1(U, x) \rightarrow \pi_1(X, x)$  trivial

then  $\pi_1(U, y) \rightarrow \pi_1(X, y)$  trivial  $\forall y \in U$  since

$$\begin{array}{ccc} \pi_1(U, x) & \rightarrow & \pi_1(X, x) \\ \phi_n \downarrow \cong & \circ & \cong \downarrow \phi_n \\ \pi_1(U, y) & \rightarrow & \pi_1(X, y) \end{array} \quad \text{h path } y \text{ to } x$$

also, if  $U \in \mathcal{U}$  and  $V \subset U$  is open and path connected

$$\text{then } \pi_1(V, x) \xrightarrow{\quad \text{trivial} \quad} \pi_1(U, x) \rightarrow \pi_1(X, x)$$

$\therefore V \in \mathcal{U}$

now  $X$  semilocally simply connected says for any  $x \in X$   
 and open set  $U$  containing  $x$ ,  $\exists$  open set  $V$  st.

$x \in V$  and  $\pi_1(V, x) \rightarrow \pi_1(X, x)$  trivial

so  $U \cap V$  open set containing  $x$  and

$$\pi_1(U \cap V, x) \rightarrow \pi_1(X, x) \text{ trivial}$$

$X$  locally path connected  $\Rightarrow \exists$  open  $W$  st.  $x \in W \subset U \cap V$  and

$W$  is path connected


from above  $\pi_1(W, x) \rightarrow \pi_1(X, x)$  trivial

so  $W \in \mathcal{U}$  and  $\mathcal{U}$  is a basis for topology on  $X$ .

for each  $U \in \mathcal{U}$  and  $\gamma$  a path  $x_0$  to a point in  $U$

set  $U_\gamma = \{ \langle \gamma * \eta \rangle \mid \eta \text{ a path in } U \text{ st. } \eta(0) = \gamma(1) \}$

note  $U_\gamma \subset \tilde{X}_H$



Claim:  $\{U_\gamma\}_{\substack{U \in \mathcal{U} \\ \gamma \text{ path } x_0 \text{ to } p \text{ in } U}}$  forms a basis for a topology on  $\tilde{X}_H$

(recall, a collection of sets in a set form a basis for a topology if given any two sets  $U, V$  in collection and a point  $x \in U \cap V$ ,  $\exists W$  in collection st.  $x \in W \subset U \cap V$  and the set is the union all elts in collection)

Pf:

note: 1) if  $\langle \gamma \rangle = \langle \delta \rangle$ , then  $U_\gamma = U_\delta$  (so can write  $U_{\langle \gamma \rangle}$ )

indeed, if  $\gamma \sim \delta$ , then  $\gamma * \eta \sim \delta * \eta \quad \forall \eta \text{ path in } U$   
H-equiv. with  $\eta(0) = \gamma(1)$


$$\text{since } (\gamma * \eta) * (\overline{\delta * \eta}) = \gamma * \eta * \bar{\eta} * \bar{\delta} \\ \simeq \gamma * \bar{\delta}$$

$$\text{so } [\gamma * \eta * \overline{\delta * \eta}] \in H.$$

2)  $p: U_{\langle \gamma \rangle} \rightarrow U$  is onto (since  $U$  path connected)

3)  $p: U_{\langle \gamma \rangle} \rightarrow U$  is one-to-one

indeed, if  $p(\langle \gamma * \eta \rangle) = p(\langle \gamma * \eta' \rangle)$ , then  $(\gamma * \eta)(1) = (\gamma * \eta')(1)$

so  $\eta * \bar{\eta}'$  is a loop in  $U$  

based at  $x \therefore \eta * \bar{\eta}'$  is homotopically trivial in  $X$

$$\Rightarrow \eta \sim \eta' \text{ rel end points}$$

$$\Rightarrow \gamma * \eta \sim \gamma * \eta' \text{ rel end points}$$

$$\Rightarrow [(\gamma * \eta) * \overline{(\gamma * \eta')}] = [e_{x_0}] \in H$$

$$\text{so } \langle \gamma * \eta \rangle = \langle \gamma * \eta' \rangle$$

4) If  $\langle \gamma' \rangle \in U_{\langle \gamma \rangle}$ , then  $U_{\langle \gamma' \rangle} = U_{\langle \gamma \rangle}$

indeed, by hypothesis  $\exists \eta$  a path in  $U$

$$\text{s.t. } \langle \gamma' \rangle = \langle \gamma * \eta \rangle$$

so we can take  $\gamma * \eta$  to represent  $\gamma'$  by 1)

if  $\langle \delta \rangle \in U_{\langle \gamma' \rangle}$ , then  $\delta = (\gamma * \eta) * \eta' = \gamma * (\eta * \eta')$

$$\text{so } \langle \delta \rangle \in U_{\langle \gamma \rangle}$$

$$\text{similarly } \langle \delta \rangle \in U_{\langle \gamma \rangle} \Rightarrow \langle \delta \rangle \in U_{\langle \gamma' \rangle}$$

now if  $\langle \delta \rangle \in U_{\langle \gamma \rangle} \cap V_{\langle \gamma' \rangle}$ , then  $U_{\langle \gamma \rangle} = U_{\langle \delta \rangle}$  and  $V_{\langle \gamma' \rangle} = V_{\langle \delta \rangle}$

so if  $W \in \mathcal{U}$  s.t.  $W \subset U \cap V$  and  $\delta(t) \in W$

then  $W_{\langle \delta \rangle} \subset U_{\langle \delta \rangle} \cap V_{\langle \delta \rangle} = U_{\langle \gamma \rangle} \cap V_{\langle \gamma' \rangle}$

clearly  $\tilde{X}_H$  is a union of all  $U_{\langle \gamma \rangle}$ 's

so we have a basis for a topology on  $\tilde{X}_H$

Claim: with above topology  $(\tilde{X}_H, \rho_H)$  is a covering space of  $X$

Pf: note:  $\forall U \in \mathcal{U}$ ,  $\delta$  paths  $x_0$  to  $pt$  in  $U$

$\rho_H|_{U_{\langle \gamma \rangle}} : U_{\langle \gamma \rangle} \rightarrow U$  a homeomorphism

indeed,  $\rho_H$  is a bijection by 2) and 3)

$\rho_H|_{U_{\langle \gamma \rangle}}$  is continuous since any basic open set  $V \subset U$   
and any path  $\delta$  from  $x_0$  to  $pt$  in  $V$

$$\left(\rho_H|_{U_{\langle \gamma \rangle}}\right)^{-1}(V) = V_{\langle \delta \rangle} \quad \therefore \text{open}$$

the above argument also shows that basic open sets in  $U_{\langle \gamma \rangle}$  map to basic open sets in  $U$

$\therefore \rho_H|_{U_{\langle \gamma \rangle}}$  a homeomorphism

note this implies  $\rho$  is continuous

now if  $x \in X$ , then let  $U \subset \mathcal{U}$  be a set containing  $x$


$\rho^{-1}(U) = \text{union of } U_{\langle \gamma \rangle}$  as  $\delta$  runs through all paths  $x_0$  to  $pt$  in  $U$

and  $\rho|_{U_{\langle \gamma \rangle}} : U_{\langle \gamma \rangle} \rightarrow U$  homeomorphism.

Claim:  $(\rho_H)_* (\pi_1(\tilde{X}_H, \tilde{x}_0)) = H$

Pf: if  $[\gamma] \in H$ , then let  $\gamma_t(s)$  be the path

$$\begin{aligned} \gamma_t : [0,1] &\rightarrow X \\ s &\longmapsto \gamma_t(s) \end{aligned}$$



note:  $\tilde{\gamma} : [0,1] \rightarrow \tilde{X}_H$  is a loop since  $\tilde{\gamma}(0) = \langle e_{x_0} \rangle$   
 $t \longmapsto \langle \gamma_t \rangle$  and  $\tilde{\gamma}(1) = \langle \gamma \rangle = \langle e_{x_0} \rangle$

since  $[\gamma] \in H$

moreover  $\rho_H \circ \tilde{\gamma} = \gamma$

$\therefore [\gamma] \in \text{image}(\rho_H)_*$


now if  $[\gamma] \notin H$ , then the path

$$\begin{aligned} \tilde{\gamma}: [0,1] &\rightarrow \tilde{X}_H \\ t &\mapsto \langle \gamma_t \rangle \end{aligned}$$

is clearly the lift of  $\gamma$  based at  $\tilde{x}_H$

and  $\tilde{\gamma}(1) = \langle \gamma \rangle$

but  $\langle \gamma \rangle \neq \langle e_{x_0} \rangle = \tilde{x}_H$  since  $[\gamma] \notin H$

$\therefore [\gamma] \notin \text{image}(\rho_H)_*$  by lemma 21,2) 


let  $p: \tilde{X} \rightarrow X$  be a covering space

a deck transformation or covering transformation is a covering space

isomorphism  $f: \tilde{X} \rightarrow \tilde{X}$

the set  $G(\tilde{X})$  of deck transformations clearly is a group under the operation of composition

examples:




$\phi_k: S' \rightarrow S'$  is a covering transformation  
 $\theta \mapsto \frac{2\pi k}{n}$

if  $f$  is any deck transformation then  $f(\tilde{x}_i) = \tilde{x}_i$  for some  $i$ , but  $\phi_i(\tilde{x}_i) = \tilde{x}_i$  too

so by lemma 24,  $f = \phi_i$  (since covering transforms are lifts of  $p_n$ )

so  $G(\tilde{X}) = \mathbb{Z}/n\mathbb{Z}$

2)



if we lift  $b$  to  $\tilde{x}_3$  then it is a loop but if we lift to  $\tilde{x}_2$  or  $\tilde{x}_1$  it is a path so no deck trans. taking  $\tilde{x}_3$  to  $\tilde{x}_1$  or  $\tilde{x}_2$  similarly can't send  $\tilde{x}_2$  to  $\tilde{x}_1$  or  $\tilde{x}_3$



so any deck trans fixes all  $\tilde{x}_i \therefore$  is identity

$$\therefore G(\tilde{X}) = \{1\}$$

A covering space  $p: \tilde{X} \rightarrow X$  is called normal if  $\forall x \in X$  and  $\tilde{x}, \tilde{x}' \in p^{-1}(x)$

$$\exists \phi \in G(\tilde{X}) \text{ s.t. } \phi(\tilde{x}) = \tilde{x}'$$

so example 1) is normal but example 2) is not.

Th<sup>m</sup> 27:

let  $p: (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$  be a path connected, locally path connected covering space of a space  $X$

let  $H = p_* (\pi_1(\tilde{X}, \tilde{x}_0)) < \pi_1(X, x_0)$ , then

1)  $(\tilde{X}, p)$  is normal  $\Leftrightarrow H$  is a normal subgroup of  $\pi_1(X, x_0)$

2)  $G(\tilde{X}) \cong N(H)/H$  where  $N(H)$  is the "normalizer" of  $H$ , i.e. largest subgroup of  $\pi_1(X, x_0)$  containing  $H$  as a normal subgroup

Remark: If  $p: \tilde{X} \rightarrow X$  normal, then  $G(\tilde{X}) = \pi_1(X, x_0) / p_* (\pi_1(\tilde{X}, \tilde{x}_0))$

in particular, for the universal cover  $p: \tilde{X} \rightarrow X$

$$G(\tilde{X}) = \pi_1(X, x_0)$$

Proof:

1)  $(\Rightarrow)$  let  $[\gamma] \in \pi_1(X, x_0)$

and  $\tilde{\gamma}$  a lift of  $\gamma$  based at  $\tilde{x}_0$

$$\text{set } \tilde{x}_1 = \tilde{\gamma}(1)$$

$$\text{from lemma 22 } [\gamma] p_* (\pi_1(\tilde{X}, \tilde{x}_1)) [\gamma]^{-1} = p_* (\pi_1(\tilde{X}, \tilde{x}_0))$$

by hypothesis  $\exists \phi \in G(\tilde{X})$  s.t.  $\phi(\tilde{x}_1) = \tilde{x}_0$

so  $\phi_*: \pi_1(\tilde{X}, \tilde{x}_1) \rightarrow \pi_1(\tilde{X}, \tilde{x}_0)$  an isomorphism

$$\therefore p_* (\pi_1(\tilde{X}, \tilde{x}_0)) = p_* \circ \phi_* (\pi_1(\tilde{X}, \tilde{x}_1)) = p_* (\pi_1(\tilde{X}, \tilde{x}_1))$$

$$\text{and } [\gamma] H [\gamma]^{-1} = H$$

$(\Leftarrow)$  let  $\tilde{x}_0$  and  $\tilde{x}_1$  be two points in  $p^{-1}(x_0)$

$$\text{and } H_1 = p_* (\pi_1(\tilde{X}, \tilde{x}_1))$$

let  $h$  be a path in  $\tilde{X}$  from  $\tilde{x}_0$  to  $\tilde{x}_i$  and  $\gamma = \rho \circ h$

by lemma 22,  $[\gamma]H[\gamma]^{-1} = H$

$\therefore H = H_i$  since  $H$  is normal

i.e.  $\rho_*(\pi_1(\tilde{X}, \tilde{x}_i)) = \rho_*(\pi_1(\tilde{X}, \tilde{x}_0))$

$\therefore$  by Th<sup>m</sup> 23  $\exists$  lifts of  $\rho$  to  $\phi_1$  and  $\phi_2$

$$\begin{array}{ccccc} (\tilde{X}, \tilde{x}_i) & \xrightarrow{\phi_1} & (\tilde{X}, \tilde{x}_0) & \xrightarrow{\phi_2} & (\tilde{X}, \tilde{x}_i) \\ & \searrow \rho & \downarrow \rho & \swarrow \rho & \\ & & (X, x_0) & & \end{array}$$

note:  $\phi_2 \circ \phi_1$  is a lift of  $\rho$  that fixes  $\tilde{x}_i$

so is  $\text{id}_{\tilde{X}}$   $\therefore \phi_2 \circ \phi_1 = \text{id}_{\tilde{X}}$

similarly  $\phi_1 \circ \phi_2 = \text{id}_{\tilde{X}}$

$\therefore \phi_1$  a deck transform taking  $\tilde{x}_i$  to  $\tilde{x}_0$

exercise: for any  $x \in X$  and  $\tilde{x}, \tilde{x}' \in \rho^{-1}(x)$  show  $\exists \phi \in G(\tilde{X})$

s.t.  $\phi(\tilde{x}) = \tilde{x}'$

2) from above if  $[\gamma]H[\gamma]^{-1} = H$  then  $\exists \phi \in G(X)$  s.t.  $\phi(\tilde{x}_0) = \tilde{\gamma}(1)$

( $\tilde{\gamma}$  lift based at  $\tilde{x}_0$ )

so we get a map


$$\Phi: N(H) \rightarrow G(\tilde{X})$$

Claim:  $\Phi$  a homomorphism

suppose  $\phi_i = \Phi([\gamma_i])$  for  $[\gamma_i] \in N(H)$   $i=1,2$

$\phi_1(\tilde{x}_0) = \tilde{x}_i$  so  $\tilde{\gamma}_1$  path  $\tilde{x}_0$  to  $\tilde{x}_i$

note:  $\tilde{\gamma}_1 * (\phi_1 \circ \tilde{\gamma}_2)$  is a path  $\tilde{x}_0$  to  $\phi_1(\tilde{x}_2)$



and  $[\rho_*(\tilde{\gamma}_1 * (\phi_1 \circ \tilde{\gamma}_2))] = [\gamma_1 * \gamma_2] = [\gamma_1] \cdot [\gamma_2]$

so  $\phi_1 \circ \phi_2$  is  $\Phi([\gamma_1] \cdot [\gamma_2])$

Claim:  $\Phi$  is surjective

let  $\phi \in G(\tilde{X})$  take a path  $h$  in  $\tilde{X}$  from  $\tilde{x}_0$  to  $\tilde{x} = \phi(\tilde{x}_0)$

$\gamma = \rho \circ h$  is a loop in  $X$

and from above  $[\gamma]H[\gamma]^{-1} = H$  so  $[\gamma] \in N(H)$


and  $\Phi([\gamma]) = \phi$

Claim:  $\ker \Phi = H$

if  $[\gamma] \in H$ , then  $\tilde{\gamma}(1) = \tilde{x}_0$  so  $\Phi([\gamma]) = id_{\tilde{x}}$

$\therefore H \subset \ker \Phi$

if  $[\gamma] \in \ker \Phi$ , then  $\tilde{\gamma}(1) = \tilde{x}_0$  and so  $\tilde{\gamma}$  a loop

$\therefore [\gamma] \in H$  

a group action on a topological space  $X$  is a pair  $(G, \rho)$  where

$G$  is a group, and

$\rho: G \rightarrow \text{Homeo}(X)$  is a homomorphism

$\uparrow$  group of homeomorphisms

if  $G$  acts on  $X$  then we can form the quotient space  $X/G$

where two points  $x_1, x_2$  are identified if  $\exists g \in G$  st.  $\rho(g)(x_1) = x_2$

this is called the orbit space

Th<sup>m</sup> 28:

let  $G$  be a group action on  $X$  such that

\*  $\forall x \in X, \exists$  a nbhd  $U$  of  $x$  so that  $g_1 U \cap g_2 U \neq \emptyset \Rightarrow g_1 = g_2$

then 1)  $p: X \rightarrow X/G$  is a normal covering space

2)  $G \cong G(X \rightarrow X/G)$  if  $X$  is path connected

3)  $G \cong \pi_1(X/G) / p_* (\pi_1(X))$  if  $X$  is path connected and locally path connected.

Proof: fairly easy


exercise or see Hatcher 

exercise: if  $G$  is finite and  $G$  acts freely on  $X$  (i.e. has no fixed points)

then the action on  $X$  satisfies \*

examples:

1)



2)  $S^n \rightarrow \mathbb{RP}^n$

$\mathbb{Z}/2\mathbb{Z}$  action