

# Math 6452 - Fall 2017

## Homework 6

Work all these problems and talk to me if you have any questions on them, but carefully write up and turn in **only** problems 3, 5, 6, 7, 9, and 10. **Due: In class on November 21.**

1. Let  $\omega$  be a 1-form on a 3-dimensional manifold  $M$ . Suppose that  $\omega$  is not zero at any point so for each  $x \in M$  the kernel  $\xi_x$  of  $\omega(x)$  is a plane in  $T_x M$ . We say that  $\xi$  is integrable if for any two vector fields  $v$  and  $w$  with values in  $\xi$  (that is  $v$  and  $w$  are sections of  $\xi$ ) we have that the Lie bracket  $[v, w]$  is also a section of  $\xi$ . For this problem assume that  $\omega$  is integrable.
  - (a) Show that  $\omega \wedge d\omega = 0$ .
  - (b) Show there exists a 1-form  $\alpha$  such that  $d\omega = \omega \wedge \alpha$ . (Hint: prove this locally and then use a partition of unity.)
  - (c) Show that  $\omega \wedge d\alpha = 0$ .
  - (d) If  $\beta$  is another 1-form such that  $d\omega = \omega \wedge \beta$  then there is a function  $f$  such that  $\beta = \alpha + f\omega$  and  $\alpha \wedge d\alpha = \beta \wedge d\beta$ .
2. Given an area form  $\omega$  on a surface  $\Sigma$  (that is a 2-form that is never zero) then one can define the divergence of a vector field  $v$  on  $\Sigma$  as the unique function  $\text{div}_\omega v$  such that

$$L_v \omega = (\text{div}_\omega v) \omega.$$

- (a) Show that if  $\omega'$  is another area form (defining the same orientation) then there is a unique positive function  $f$  such that  $\omega' = f\omega$  and that

$$\text{div}_\omega(v) = \text{div}_{\omega'}(v) + d(\ln f)(v).$$

- (b) Derive a formula for  $\text{div}_\omega(v')$  in terms of  $\text{div}_\omega(v)$  if  $v' = gv$  for some function  $g$ .
  - (c) Show that given a function  $f : \Sigma \rightarrow \mathbb{R}$  there is a unique vector field  $v_f$  that satisfies  $\iota_{v_f} \omega = df$ .
  - (d) Show the flow of  $v_f$  from the previous item preserves the level sets of  $f$  and has zero divergence.
3. Let  $a : S^n \rightarrow S^n$  be the antipodal map, that is the map  $a(x) = -x$  when we think of  $S^n$  as the unit sphere in  $\mathbb{R}^n$ . Show that  $a$  is orientation preserving if and only if  $n$  is odd.
  4. Show that  $\mathbb{R}P^n$  is orientable if and only if  $n$  is odd.
  5. Suppose that  $M$  and  $N$  are oriented manifolds and  $f : M \rightarrow N$  is a local diffeomorphism. If  $M$  is connected then show that  $f$  is either orientation preserving or orientation reversing.
  6. On  $\mathbb{R}^n - \{0\}$  consider the  $(n-1)$ -form

$$\omega = \frac{1}{\|x\|^n} \sum_{i=1}^n (-1)^{i-1} x^i dx^1 \wedge \dots \wedge \widehat{dx^i} \wedge \dots \wedge dx^n.$$

Compute  $d\omega$ .

7. Let  $S^2$  be the unit sphere in  $\mathbb{R}^3$  and  $\omega$  the 2-form from the previous exercise. If  $i : S^2 \rightarrow \mathbb{R}^3$  is the inclusion map then compute

$$\int_{S^2} i^* \omega.$$

Is there a 1-form  $\eta$  on  $\mathbb{R}^3 - \{0\}$  such that  $d\eta = \omega$ ? Explain why or why not. Notice that this and the previous exercise imply that  $H_{DR}^2(\mathbb{R}^3 - \{0\}) \neq 0$ .

If you feel like it maybe try to work this problem again for  $S^{n-1}$  (this is not required to be turned in).

8. Use Stokes theorem to prove the classical Green's formula: Give a region  $R$  in  $\mathbb{R}^2$  with smooth boundary  $\partial R = \gamma$  then show

$$\int_{\gamma} f dx + g dy = \int_R \left( \frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dx dy.$$

9. Given any embedding  $f : T^2 \rightarrow S^3$  show that for any closed 2-form  $\omega$  on  $S^3$  we have

$$\int_{T^2} f^* \omega = 0.$$

Hint: Show that there is a smooth homotopy  $H : T^2 \times [0, 1] \rightarrow S^3$  from  $f$  to a constant map. Now use Stokes theorem.

10. Show there is some embedding  $f : T^2 \rightarrow T^3$  and a closed 2-form  $\omega$  on  $T^3$  such that

$$\int_{T^2} f^* \omega \neq 0.$$

Notice that this problem together with the previous one implies that  $S^3$  is not diffeomorphic to  $T^3$ .